

麦類赤かび病の多発条件出現日(アメダス実測値)

2026/5/19 現在

	本年 (R8)			前年 (R7)			多発年 (R6)		
	熊本市	菊池市	人吉市	熊本市	菊池市	人吉市	熊本市	菊池市	人吉市
4/1							●		◎
4/2	○		○				◎	○	○
4/3							●	●	◎
4/4	◎	◎	◎				◎	◎	◎
4/5	◎	○	○	○			◎	◎	◎
4/6				○	○	○	◎	◎	◎
4/7		○	○	○			◎	◎	◎
4/8							●	◎	◎
4/9	◎	○					◎	◎	◎
4/10	●	●	●	●	◎	◎	○	○	○
4/11	●	◎	●	◎	○	◎			
4/12	●	◎	◎	○	○	○			
4/13	●	◎	◎	○					◎
4/14	●	◎	◎	○					○
4/15	◎	◎	●				●	●	◎
4/16	●	●	●	○	○	○	●	●	●
4/17									
4/18	●	●	●	●	●	●			◎
4/19	●	●	◎	●	●	●			○
4/20	●	●	●	●	●	●	◎	◎	◎
4/21		◎	●	●	●	●	●	●	●
4/22	◎	◎	◎	●	●	●	◎	◎	●
4/23	◎	◎	◎	◎	◎	●	●	●	●
4/24	◎	◎	◎	◎	◎	◎	●	◎	●
4/25	●	◎	◎				◎	○	
4/26	◎	◎	◎				◎		◎
4/27	●	◎	◎				●	●	●
4/28				◎	◎	◎	●	●	●
4/29	◎	◎	●	◎	○	○	●	●	●
4/30	◎	◎	◎				●	●	●
5/1	◎	◎	◎	●	●	◎	◎	◎	●
5/2	●	○	◎	◎	◎	◎	◎	○	●
5/3	◎	◎	◎	●	○	○			
5/4	◎	◎	◎						
5/5				○	○	○			
5/6				◎	◎	◎	●	●	●
5/7				●	◎	●	●	●	●
5/8							◎	○	
5/9				●	●	◎	○	○	
5/10				●	●	●			
5/11				◎	◎	◎			
5/12	●	●	●	◎	◎	◎	●	●	●
5/13	●	●	●				●	●	●
5/14	●	●	●				●	●	●
5/15	●	●	●						
5/16				●	●	●			
5/17				●	●	●			
発病穂率(%) 5月上中旬調査	小麦: 13.3 大麦: 2.7			小麦: 3.8 大麦: 0.8			小麦: 24.1 大麦: 1.8		

【発生予察事業調査実施基準より】

- : 子のう殻形成の好適条件 (日平均気温13℃以上、当日または前日に降雨があった)
- ◎: 子のう胞子飛散の好適条件 (日最高気温15℃以上、日最低気温10℃以上、当日または前日に降雨があった)
- : 多発条件 (日平均気温が18℃以上、当日または前日に降雨があった)

<防除適期> 小麦 1回目:穂揃い期の10日後(出穂期12日~14日後)、2回目:1回目の7~10日後
 小麦 1回目:開花始~開花最盛期(出穂期7日~10日後)、2回目:1回目の7日~10日後